

Rx

Recombinant human Epidermal Growth Factor gel

REGEN-D® 60

For Quick Burn Healing

1. NAME AND DESCRIPTION OF THE MEDICINAL PRODUCT

REGEN-D® 60 contains human epidermal growth factor-based gel produced by recombinant DNA technology and developed by Bharat Biotech International Limited, Hyderabad, India. The primary structure of recombinant human EGF is a single chain polypeptide which is a 53 amino acids chain.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each gram of gel contains

Purified Bulk of rh - Epidermal Growth Factor: 60µg
Sodium Methylparaben IP 1.8mg
Sodium Propylparaben IP 0.2mg
Excipients q.s

3. PHARMACEUTICAL FORM

Gel.

4. CLINICAL PARTICULARS

4.1 Therapeutic Indications

REGEN-D® 60 is indicated for first and second-degree burn wounds, for healing of donor site skin graft area.

4.2 Posology, Schedule and Method of Administration

Cleanse the burn wound and surrounding surface with water or saline and pat dry with sterile cotton before the gel is applied. Apply the gel evenly (topical application) on the affected area of the skin using sterile cotton swab twice a day till the wound area heals. **REGEN-D® 60** therapy should be continued up to a period of 2 to 3 weeks after the wound heals. The continuation of the therapy is at the discretion of the physician.

- A single tube should be used for individual patient.
- Avoid direct contact of the tube with the wound area

Any unwarranted use of the product is not the responsibility of the manufacturer.

4.3 Contraindications

REGEN-D® 60 is generally well tolerated. However, the product should not be applied or repeated to persons known to be hypersensitive to any of the components of the product. Also, it should not be applied to individuals who are receiving immunosuppressive or immune-stimulant therapy, or in immune compromised individuals.

4.4 Special Warnings and Precautions

It is suggested that the medical practitioner ascertain the hypersensitivity status of the subject.

4.5 Interactions with Other Medicinal Products

REGEN-D® 60 must not be used with other growth factor containing gel or cream.

4.6 Pregnancy and Lactation:

REGEN-D® 60 is contraindicated for use in pregnant and lactating woman.

4.7 Effects on Ability to Drive and Use Machines

Since the product is for topical application, systemic absorption is not expected. However, no studies on the effect of **REGEN-D® 60** on the ability to drive and use machines has been performed.

4.8 Undesirable Effects

REGEN-D® 60 has proven low reactivity and is well tolerated, however, Skin irritation/pain, rash at the application site may be seen in very few cases.

4.9 Over-dose

Not applicable.

4.10 Pre-Clinical & Clinical Trial Experience

Pre-clinical toxicological studies done on rats and rabbits concluded that the rh-EGF is safe and well tolerable with no systemic observations. The study was conducted to evaluate the potential toxicity of repeated doses (75-300 µg/Kg) of recombinant human Epidermal Growth Factor applied topically to rats and New Zealand white rabbits. groups. The rh-EGF was not absorbed systemically as revealed in the systemic absorption study conducted in rabbit. There has been significant increase in the DNA and collagen contents in the skin samples treated with rh-EGF. No significant changes were observed control and treated groups with respect to protein contents in the skin.

In another study with rats (Wistarfurth) and rabbits (New Zealand white) to evaluate the potential toxicity of repeated doses (75-300 µg/Kg) of rh-EGF applied topically to rats and rabbits. It was found that there was no observable antibody response in the treated groups with EGF in both rats and rabbits. The DNA content in skin sample of treated group has significantly increased in high dose in both the species on 15th and 31st day. The protein content in the control and treated groups did not differ significantly in both the species studied. The collagen content was significantly increased in medium and high dose groups in males and females both the species on 15th and 31st days.

In another study, the application on burns have reduced the healing time to an average of 9 days compared to the usual healing time of 20 days. There were no cases of adverse or serious adverse events observed during the study. Only one case of rash was observed during the study, which after laboratory investigations, confirmed that this was not related to study drug medication. The laboratory investigations showed no difference in the pre- and post-drug administration values.

A multi-centre, double-blind, randomised, parallel, phase 3 study was conducted to evaluate the safety and efficacy of **REGEN-D® 60** in comparison with silver sulfadiazine as a treatment for donor site graft and burns. The study showed that application of **REGEN-D® 60** on donor site grafts have reduced the healing time to an average of 9-10 days compared to the regular healing time of 16-20 days.

4.11 Post-Marketing Experience

More than a hundred thousand tubes of various dose strengths have been sold in domestic and international markets since the market authorization. The product is under close surveillance for its safety in the field.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic Properties

EGF is part of a complex network of growth factors and receptors that together help to modulate the growth of cells. EGF is released by cells, and then is picked up either by the cell itself, stimulating its own growth, or by neighboring cells, stimulating their ability to divide. Receptors on the surface of the cell bind to EGF and relay the signal inside. When the receptor binds to EGF, it is activated by forming a dimer with other receptors.

EGF is essential for mediating the de-differentiation of keratinocytes to an epithelial lineage and to reestablish the epithelial barrier. EGF binds to the EGFR, a protein tyrosine kinase receptor, expressed on the majority of cells in the skin. Activation of EGFR leads to a number of biological responses, including migration, proliferation, cytoprotection, cellular differentiation, and apoptosis. In wound healing EGFR plays an important role in re-epithelialization and dermal maturation. Topical use of recombinant human EGF has been shown to increase re-epithelialization and enhance wound healing.

5.2 Pharmacokinetic Properties

Subjects were followed up for various periods of time to evaluate the systemic absorption of **REGEN-D® 60** in blood. Sera was analyzed for anti-EGF titers by Indirect ELISA method. The test serum absorbance was less than the seroconversion cut-off value, hence these samples were negative for anti r-human EGF antibody.

Patients with wounds were tested for the presence of rhEGF by collecting the samples from the site of application, the result clearly shows that rhEGF is available at the site of application. Protease enzyme present in the body degrades rhEGF at the site of applications, however when **REGEN-D® 60** was applied, there was sufficient high concentration of rhEGF locally.

6. PHARMACEUTICAL PARTICULARS

Category: Growth factor.

6.1 List of Excipients

Sodium Methylparaben IP
Sodium Propylparaben IP

6.2 Incompatibilities

Recombinant human EGF is combined along with silver sulfadiazine and chlorhexidine, marketed as **SLVRGEN®** is manufactured by Bharat Biotech and is used for the treatment of first and second-degree burns and ulcers like abrasions, incisions, minor cuts and wounds. No other studies were performed to look for incompatibilities with **REGEN-D® 60**.

6.3 Shelf Life

The expiry date of the product is indicated on the carton.

6.4 Storage

Store in a cool and dry place. Do not freeze. Do not use after expiry date.

7. PRESENTATION

REGEN-D® 60 is presented in aluminum tube with polypropylene screw cap. The pack sizes are: 7.5gm, 15gm, 30gm, 60gm and 150gm.

Last revision date: September 2019

Manufactured by:

∞∞
BHARAT
BIOTECH
Lead Innovation

Bharat Biotech International Ltd.

Genome Valley, Shameerpet Mandal,
Medchal-Malkajgiri District-500078, Telangana, India.

For complaints and suggestions about the product, and any adverse event,
Please email feedback@bharatbiotech.com or call on Toll free number 1800 102 2245

2301000041
180920